

Measures of closeness to cordiality for graphs

Summary of Results

Anand Brahmbhatt* Kartikeya Rai† Amitabha Tripathi‡§

Abstract

Keywords. cordial labelling
2010 MSC.

1 Introduction

Let $f : V(G) \rightarrow \{0, 1\}$, and let $\bar{f} : E(G) \rightarrow \{0, 1\}$ be the induced mapping given by

$$\bar{f}(xy) = |f(x) - f(y)|.$$

We say that f is a *cordial* labelling of G if the number of vertices labelled 0 and the number of vertices labelled 1 differ by at most 1, and the number of edges labelled 0 and the number of edges labelled 1 differ at most by 1.

For $i \in \{0, 1\}$, let $v_i(f)$ denote the number of vertices labelled i and let $e_i(f)$ denote the number of edges labelled i . Let

$$\Delta_v(f) = |v_0(f) - v_1(f)| \text{ and } \Delta_e(f) = |e_0(f) - e_1(f)|.$$

We introduce two measures of closeness to cordiality of a graph.

$$\mathcal{D}_1(G) = \min \left\{ \Delta_v(f) + \Delta_e(f) \mid f : V(G) \rightarrow \{0, 1\} \right\}. \quad (1)$$

$$\mathcal{D}_2(G) = \min \left\{ \Delta_e(f) \mid f : V(G) \rightarrow \{0, 1\}, \Delta_v(f) \leq 1 \right\}. \quad (2)$$

*Department of Computer Science & Engineering, Indian Institute of Technology, Hauz Khas, New Delhi – 110016, India.

e-mail:anand.brahmbhatt@iitdalumni.com

†Department of Mathematics, Indian Institute of Technology, Hauz Khas, New Delhi – 110016, India.
e-mail:mt6180811@iitd.ac.in

‡Department of Mathematics, Indian Institute of Technology, Hauz Khas, New Delhi – 110016, India.
e-mail:atripath@maths.iitd.ac.in

§Corresponding author

2 The Tree

Theorem 2.1. Let \mathcal{T}_n denote a tree with n vertices.

$$\mathcal{D}_1(\mathcal{T}_n) = 1 \quad \text{and} \quad \mathcal{D}_2(\mathcal{T}_n) = \begin{cases} 0 & \text{if } n \equiv 1 \pmod{2} \\ 1 & \text{if } n \equiv 0 \pmod{2} \end{cases}$$

3 The complete graphs \mathcal{K}_n

Theorem 3.1. If $a^2 \leq n < (a+1)^2$, then

$$\mathcal{D}_1(\mathcal{K}_n) = \begin{cases} a + \frac{1}{2}(n - a^2) & \text{if } n \in \{a^2 + 2t : 0 \leq t \leq a\}; \\ 2a - 1 & \text{if } n = a^2 + 1; \\ a + 1 + \frac{1}{2}((a+1)^2 - n) & \text{if } n \in \{a^2 + 2t + 1 : 1 \leq t \leq a-1\}. \end{cases}$$

$$\mathcal{D}_2(\mathcal{K}_n) = \left\lfloor \frac{n}{2} \right\rfloor.$$

4 The complete r -partite graphs $\mathcal{K}_{n_1, \dots, n_r}$

Theorem 4.1. If n_1, \dots, n_r are positive integers of which s are odd, then

$$\mathcal{D}_1(\mathcal{K}_{n_1, \dots, n_r}) \leq \begin{cases} \frac{1}{2} \left(s + 1 - (2a-1)^2 \right) & \text{if } (2a)^2 \leq s \leq (2a+1)^2 \text{ and } s \equiv 0 \pmod{2} \\ \frac{1}{2} \left(s + 1 - (2a-2)^2 \right) & \text{if } (2a)^2 \leq s \leq (2a+1)^2 \text{ and } s \equiv 1 \pmod{2} \\ \frac{1}{2} \left(s + 1 - (2a-1)^2 \right) & \text{if } (2a+1)^2 \leq s \leq (2a+2)^2 \text{ and } s \equiv 0 \pmod{2} \\ \frac{1}{2} (s + 1 - 4a^2) & \text{if } (2a+1)^2 \leq s \leq (2a+2)^2 \text{ and } s \equiv 1 \pmod{2} \end{cases}$$

$$\mathcal{D}_2(\mathcal{K}_{n_1, \dots, n_r}) = \left\lfloor \frac{s}{2} \right\rfloor.$$

5 The cycles \mathcal{C}_n

Theorem 5.1.

$$\mathcal{D}_1(\mathcal{C}_n) = \begin{cases} 0 & \text{if } n \equiv 0 \pmod{4}; \\ 2 & \text{if } n \not\equiv 0 \pmod{4}. \end{cases} \quad \text{and} \quad \mathcal{D}_2(\mathcal{C}_n) = \begin{cases} 0 & \text{if } n \equiv 0 \pmod{4}; \\ 1 & \text{if } n \equiv 1, 3 \pmod{4}; \\ 2 & \text{if } n \equiv 2 \pmod{4}. \end{cases}$$

Corollary 5.1. An Eulerian graph G with $4k+2$ edges is not cordial for all $k \in \mathbb{Z}^+$.

6 The wheel graphs \mathcal{W}_n

Theorem 6.1.

$$\mathcal{D}_1(\mathcal{W}_n) = \begin{cases} 0 & \text{if } n \equiv 2 \pmod{4}; \\ 1 & \text{if } n \equiv 1, 3 \pmod{4}; \\ 2 & \text{if } n \equiv 0 \pmod{4}. \end{cases} \quad \text{and} \quad \mathcal{D}_2(\mathcal{W}_n) = \begin{cases} 0 & \text{if } n \not\equiv 0 \pmod{4}; \\ 2 & \text{if } n \equiv 0 \pmod{4}. \end{cases}$$

7 The fan graphs $\mathcal{F}_{m,n}$

Theorem 7.1.

$$\mathcal{D}_1(\mathcal{F}_{m,n}) = \begin{cases} 1 & \text{if } n \equiv 0, 2 \pmod{4} \text{ and } m \equiv 0, 2 \pmod{4}; \\ 2 & \text{if } n \equiv 0, 2 \pmod{4} \text{ and } m \equiv 1, 3 \pmod{4}; \\ 1 & \text{if } n \equiv 1, 3 \pmod{4} \text{ and } m \equiv 0, 2 \pmod{4}; \\ 1 & \text{if } n \equiv 1, 3 \pmod{4} \text{ and } m \equiv 1, 3 \pmod{4}. \end{cases} \quad (3)$$

$$\mathcal{D}_2(\mathcal{F}_{m,n}) = \begin{cases} 1 & \text{if } n \equiv 0, 2 \pmod{4} \text{ and } m \equiv 0, 2 \pmod{4}; \\ 1 & \text{if } n \equiv 0, 2 \pmod{4} \text{ and } m \equiv 1, 3 \pmod{4}; \\ 0 & \text{if } n \equiv 1, 3 \pmod{4} \text{ and } m \equiv 0, 2 \pmod{4}; \\ 1 & \text{if } n \equiv 1, 3 \pmod{4} \text{ and } m \equiv 1, 3 \pmod{4}. \end{cases} \quad (4)$$

8 Bound on $\mathcal{D}_1(G_1 \vee G_2)$ and $\mathcal{D}_2(G_1 \vee G_2)$

Theorem 8.1. $G_1 \vee G_2$ is a graph join between G_1 and G_2 . It is defined as $G_1 \vee G_2 = (V(G_1) \cup V(G_2), E(G_1) \cup E(G_2) \cup V(G_1) \times V(G_2))$. Then,

$$\mathcal{D}_1(G_1 \vee G_2) \leq \mathcal{D}_1(G_1) + \mathcal{D}_1(G_2) + \mathcal{D}_1(G_1)\mathcal{D}_1(G_2) \quad (5)$$

$$\mathcal{D}_2(G_1 \vee G_2) \leq \mathcal{D}_2(G_1) + \mathcal{D}_2(G_2) + 1 \quad (6)$$